

BL21 感受态细胞

BL21 Chemically Competent Cell 说明书

产品货号: ML-G2024

保存条件: -80℃

产品规格: 10×100μl 50×100μl

产品介绍

基 因 型

E. coli B F- dcm ompT hsdS(rB- mB-) gal [malB+]K-12(λ S)

简 要 说 明

BL21 是最早开发的用于原核表达的菌株，BL21（DE3）、Rosetta、

OrigamiB(DE3) 等一系列原核表达菌株均来源于 BL21 菌株。该菌株主要用于非毒性蛋白的表达，不含 T7 RNA 聚合酶，所以不能用于由 T7 启动子驱动蛋白表达(如：pET 系列)；但含有大肠杆菌 RNA 聚合酶，可以用于 tac 或 trc 等使用大肠杆菌 RNA 聚合酶的原核系统的表达（如：pGEX，pMAL 质粒）。MLBio High5TM 系列 BL21 感受态细胞由特殊工艺制作，经 pUC19 质粒检测转化效率达 107cfu/ μ g。

操 作 说 明

1. 取 100 μ l 冰上融化的 BL21 感受态细胞，加入目的质粒并轻轻混匀，冰上静置 25 分钟。
2. 42℃ 水浴热激 45 秒，迅速放回冰上并静置 2 分钟，晃动会降低转化效率。
3. 向离心管中加入 700 μ l 不含抗生素的无菌培养基(2YT 或 LB)，混匀后 37℃，200rpm 复苏 60 分钟。
4. 5000rpm 离心一分钟收菌，留取 100 μ l 左右上清轻轻吹打重悬菌块并涂布到含相应抗生素的 2YT 或 LB 培养基上。
5. 将平板倒置放于 37℃ 培养箱过夜培养。

注 意 事 项

1. MLBio 感受态细胞最好在冰上融化。

2. 混入质粒时应轻柔操作。
3. 转化高浓度的质粒可相应减少最终用于涂板的菌量。
4. 诱导时，IPTG 浓度可选(0.1-2mM 均可)。
5. 为获得需要量的蛋白，最佳诱导时间，温度，IPTG 浓度需实验者优化。

Sample Induction Protocol (for reference only)

1. Inoculate a single colony from a freshly streaked plate into 3ml of LB medium containing the appropriate antibiotic for the plasmid and host strain.
2. Incubate with shaking at 200 rpm at 37°C overnight.
3. Inoculate 50 ml of LB medium containing the appropriate antibiotic with 0.5 ml of the overnight culture prepared in step 2(use the 500 ml triangular flask as the container would be better).
4. Incubate with shaking at 150 rpm at 37°C until the OD 600 reaches 0.5-0.8. (0.6 recommended; about 2.5h).
5. (Optional)Pipet 1ml of the cultures into clean microcentrifuge tubes and place the tubes on ice until needed for gel analysis or storage at -20°C. These will serve as the non-induced control samples.
6. Add IPTG to a final concentration of 1 mM. Optimal time for induction of the target protein may vary from 2-16 hours, depending on the protein.
7. Incubate with shaking at 120 rpm at 37°C for 2-4 hours. To determine the optimal time for induction of the target protein, it is recommended that a time course experiment be performed varying the induction from 2-16 hours.
8. Place the culture on ice for 10 minutes. Harvest cells by centrifugation at 5,000 x g for 10 minutes at 4°C.
9. Remove the supernatant and store the cell pellet at -20°C (storage at lower temperatures is also acceptable).

IPTG

Prepare a 1 M solution of IPTG (Isopropyl- β -D-thiogalactoside; Isopropyl- β -D-thiogalactopyranoside) by dissolving 2.38 g of IPTG in dd water and adjust the final volume to 10 ml. Filter sterilize before use.