

兔抗 ABL1/ABL2(phospho-Tyr393/439)多克隆抗体

中文名称：兔抗 ABL1/ABL2 (phospho-Tyr393/439)多克隆抗体

英文名称：Anti-ABL1/ABL2 (phospho-Tyr393/439) rabbit polyclonal antibody

别名：Abelson murine leukemia viral oncogene homolog 1; c-ABL; p150

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：ABL1/ABL2(phospho-Tyr393/439)

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

Regulates cytoskeleton remodeling during cell differentiation, cell division and cell adhesion. Localizes to dynamic actin structures, and phosphorylates CRK and CRKL, DOK1, and other proteins controlling cytoskeleton dynamics. Regulates DNA repair potentially by activating the proapoptotic pathway when the DNA damage is too severe to be repaired. Phosphorylates PSMA7 that leads to an inhibition of proteasomal activity and cell cycle tr

	ansition blocks.
Applications:	WB, IHC, IF
Name of antibody:	ABL1/ABL2(phospho-Tyr393/439)
Immunogen:	Peptide sequence around phosphorylation site of tyrosine 393/439 (D-T-Y(p)-T-A) derived from Human ABL1/2.
Full name:	ABL proto-oncogene 1, non-receptor tyrosine kinase/ABL proto-oncogene 2, non-receptor tyrosine kinase
Synonyms:	ABL; JTK7; p150; c-ABL; v-abl; CHDSKM; c-ABL1; bcr/abl; ARG; ABLL
SwissProt:	P00519/P42684
IHC positive control:	Human breast carcinoma
IHC Recommend dilution:	50-100
WB Predicted band size:	210 kDa
WB Positive control:	HL60 cells
WB Recommended dilution:	500-1000
IF Positive control:	Hela cells
IF Recommended dilution:	100-200





