

兔抗 CHST13 多克隆抗体

中文名称：兔抗 CHST13 多克隆抗体

英文名称： Anti-CHST13 rabbit polyclonal antibody

别名： C4ST3

抗原： CHST13

储存： 冷冻（-20℃） 避光

宿主： Rabbit

反应种属： Human

相关类别： 一抗

标记物： Unconjugate

克隆类型： Unconjugate

技术规格

| | |
|--------------------------|---|
| Background: | The protein encoded by this gene belongs to the sulfotransferase 2 family. It is localized to the golgi membrane, and catalyzes the transfer of sulfate to the C4 hydroxyl of beta -1,4-linked N-acetylgalactosamine (GalNAc) flanked by glucuronic acid residue in chondroitin. Chondroitin sulfate constitutes the predominant proteoglycan present in cartilage and is distributed on the surfaces of many cells and extracellular matrices. |
| Applications: | WB, IHC |
| Name of antibody: | CHST13 |
| Immunogen: | Synthesized peptide derived from internal of human CHST1 |

| | |
|---------------------------------|--|
| | 3. |
| Full name: | carbohydrate (chondroitin 4) sulfotransferase 13 |
| Synonyms : | C4ST3 |
| SwissProt: | Q8NET6 |
| IHC positive control: | Human testis tissue |
| IHC Recommend dilution: | 50-100 |
| WB Predicted band size: | 39 kDa |
| WB Positive control: | K562 cells lysate |
| WB Recommended dilution: | 500-3000 |



