

## 兔抗 ASCC1 多克隆抗体

中文名称：兔抗 ASCC1 多克隆抗体

英文名称：Anti-ASCC1 rabbit polyclonal antibody

别名：p50; CGI-18; ASC1p50

相关类别：一抗

抗原：ASCC1

储存：冷冻（-20℃）

宿主：Rabbit

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

#### Background:

This gene encodes a subunit of the activating signal cointegrator 1 (ASC-1) complex. The ASC-1 complex is a transcriptional coactivator that plays an important role in gene transactivation by multiple transcription factors including activating protein 1 (AP-1), nuclear factor kappa-B (NF-kB) and serum response factor (SRF). The encoded protein contains an N-terminal KH-type RNA-binding motif which is required for AP-1 transactivation by the ASC-1 complex. Mutations in this gene are associated with Barrett esophagus and esophageal adenocarcinoma

	. Alternatively spliced transcripts encoding multiple isoforms have been observed for this gene.
<b>Applications:</b>	ELISA, IHC
<b>Name of antibody:</b>	ASCC1
<b>Immunogen:</b>	Fusion protein of human ASCC1
<b>Full name:</b>	activating signal cointegrator 1 complex subunit 1
<b>Synonyms :</b>	p50; CGI-18; ASC1p50
<b>SwissProt:</b>	Q8N9N2
<b>ELISA Recommended dilution:</b>	2000-5000
<b>IHC positive control:</b>	Human thyroid cancer and human esophagus cancer
<b>IHC Recommend dilution:</b>	25-100

