

## 兔抗 CYP2C9 多克隆抗体

中文名称：兔抗 CYP2C9 多克隆抗体

英文名称： Anti-CYP2C9 rabbit polyclonal antibody

别名： CPC9; CYP2C; CYP2C10; CYP2C9; P450IIC9

相关类别： 一抗

抗原： CYP2C9

储存： 冷冻 (-20℃)

宿主： Rabbit

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格	This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by rifampin. The enzyme is known to metabolize many xenobiotics, including phenytoin, tolbutamide, ibuprofen and S-warfarin. Studies identifying individuals who are poor metabolizers of phenytoin and tolbutamide suggest that this gene is polymorphic. The gene is located within a cluster of cytochrome P450 genes on chromosome 10q24.
<b>Background:</b>	
<b>Applications:</b>	ELISA, IHC
<b>Name of antibody:</b>	CYP2C9
<b>Immunogen:</b>	Fusion protein of human CYP2C9

<b>Full name:</b>	cytochrome P450, family 2, subfamily C, polypeptide 9
<b>Synonyms :</b>	CPC9; CYP2C; CYP2C10; CYP2C9; P450IIC9
<b>SwissProt:</b>	P11712
<b>ELISA Recommended dilution:</b>	2000-5000
<b>IHC positive control:</b>	Human liver cancer
<b>IHC Recommend dilution:</b>	50-200

