

兔抗 BMX(Phospho-Tyr566) 多克隆抗体

中文名称: 兔抗 BMX(Phospho-Tyr566) 多克隆抗体

英文名称: Anti-BMX(Phospho-Tyr566) rabbit polyclonal antibody

别 名: ETK; PSCTK2; PSCTK3

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

抗 原: BMX(Phospho-Tyr566)

反应种属: Human Mouse

标记物: Unconjugate

克隆类型: Unconjugate

技术规格

	This gene encodes a non-receptor tyrosine kinase belongi
	ng to the Tec kinase family. The protein contains a PH-lik
	e domain, which mediates membrane targeting by binding
	to phosphatidylinositol 3,4,5-triphosphate (PIP3), and a SH
Background:	2 domain that binds to tyrosine-phosphorylated proteins a
	nd functions in signal transduction. The protein is implicat
	ed in several signal transduction pathways including the St
	at pathway, and regulates differentiation and tumorigenicit
	y of several types of cancer cells.
Applications:	WB
Name of antibody:	BMX(Phospho-Tyr566)



Immunogen:	Peptide sequence around phosphorylation site of Tyrosine 566 (D-Q-Y(p)-V-S) derived from Human ETK.
Full name:	BMX non-receptor tyrosine kinase
Synonyms:	ETK; PSCTK2; PSCTK3
SwissProt:	P51813
WB Predicted band size:	78 kDa
WB Positive control:	Hela cells lysates treated with Serum
WB Recommended dilution:	500-1000

