

## 兔抗 CD2AP 多克隆抗体

中文名称: 兔抗 CD2AP 多克隆抗体

英文名称: Anti-CD2AP rabbit polyclonal antibody

别 名: CMS

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: CD2AP

反应种属: Human

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

**Background:** 

This gene encodes a scaffolding molecule that regula tes the actin cytoskeleton. The protein directly interac ts with filamentous actin and a variety of cell membr ane proteins through multiple actin binding sites, SH 3 domains, and a proline-rich region containing binding sites for SH3 domains. The cytoplasmic protein lo calizes to membrane ruffles, lipid rafts, and the leading edges of cells. It is implicated in dynamic actin re modeling and membrane trafficking that occurs during receptor endocytosis and cytokinesis. Haploinsufficiency of this gene is implicated in susceptibility to glo



|                             | merular disease.                      |
|-----------------------------|---------------------------------------|
| Applications:               | ELISA, WB, IHC                        |
| Name of antibody:           | CD2AP                                 |
| Immunogen:                  | Synthetic peptide of human CD2AP      |
| Full name:                  | CD2-associated protein                |
| Synonyms:                   | CMS                                   |
| SwissProt:                  | Q9Y5K6                                |
| ELISA Recommended dilution: | 2000-10000                            |
| IHC positive control:       | Human Lymphoma and Human liver cancer |
| IHC Recommend dilution:     | 100-300                               |
| WB Predicted band size:     | 71 kDa                                |
| WB Positive control:        | A431 and Raji cells                   |
| WB Recommended dilution:    | 1000-5000                             |





