

## 兔抗 CD22(Phospho-Tyr807) 多克隆抗体

中文名称：兔抗 CD22(Phospho-Tyr807) 多克隆抗体

英文名称：Anti-CD22(Phospho-Tyr807) rabbit polyclonal antibody

别名：SIGLEC2; SIGLEC-2

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：CD22(Phospho-Tyr807)

反应种属：Human

标记物：Unconjugate

克隆类型：Unconjugate

### 技术规格

<b>Background:</b>	Mediates B-cell B-cell interactions. May be involved in the localization of B-cells in lymphoid tissues. Binds sialylated glycoproteins; one of which is CD45. Preferentially binds to alpha2,6-linked sialic acid. The sialic acid recognition site can be masked by cis interactions with sialic acids on the same cell surface. Upon ligand induced tyrosine phosphorylation in the immune response seems to be involved in regulation of B cell antigen receptor signaling. Plays a role in positive regulation through interaction with Src family tyrosine kinases and may also act as an inhibitory receptor by recruiting
--------------------	---

	cytoplasmic phosphatases via their SH2 domains that block signal transduction through dephosphorylation of signaling molecules.
<b>Applications:</b>	WB
<b>Name of antibody:</b>	CD22(Phospho-Tyr807)
<b>Immunogen:</b>	Peptide sequence around phosphorylation site of tyrosine 807 (G-D-Y(p)-E-N) derived from Human BL-CAM.
<b>Full name:</b>	CD22 molecule
<b>Synonyms :</b>	SIGLEC2; SIGLEC-2
<b>SwissProt:</b>	P20273
<b>WB Predicted band size:</b>	95 kDa
<b>WB Positive control:</b>	COS7 cells lysates
<b>WB Recommended dilution:</b>	500-1000

