

兔抗 EIF4G1 (phospho-Ser1232)多克隆抗体

中文名称：兔抗 EIF4G1 (phospho-Ser1232)多克隆抗体

英文名称：Anti-EIF4G1 (phospho-Ser1232) rabbit polyclonal antibody

别名：P220; EIF4F; EIF4G; EIF4GI; PARK18; EIF-4G1

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：EIF4G1 (phospho-Ser1232)

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	eIF4F is a multi-subunit complex, the composition of which varies with external and internal environmental conditions. It is composed of at least EIF4A, EIF4E and EIF4G1/EIF4G3. Interacts with eIF3, mutually exclusive with EIF4A1 or EIFA2, EIF4E and through its N-terminus with PAPB C1. Interacts through its C-terminus with the serine/threonine kinases MKNK1, and with MKNK2. Appears to act as a scaffold protein, holding these enzymes in place to phosphorylate EIF4E. Non-phosphorylated EIF4EBP1 competes with EIF4G1/EIF4G3 to interact with EIF4E; insulin stimulated MAP-kinase (MAPK1 and MAPK3) phosphorylates
--------------------	--

	on of EIF4EBP1 causes dissociation of the complex allowing EIF4G1/EIF4G3 to bind and consequent initiation of translation. EIF4G1/EIF4G3 interacts with PABPC1 to bring about circularization of the mRNA. Rapamycin can attenuate insulin stimulation mediated by FKBP. Interacts with EIF4E3. Interacts with MIF4GD. Interacts with rotavirus A NSP3; in this interaction, NSP3 takes the place of PABPC1 thereby inducing shutoff of host protein synthesis
Applications:	WB, IHC
Name of antibody:	EIF4G1 (phospho-Ser1232)
Immunogen:	Synthetic peptide of human EIF4G1 (phospho-Ser1232)
Full name:	eukaryotic translation initiation factor 4 gamma, 1 (phospho-Ser1232)
Synonyms :	P220; EIF4F; EIF4G; EIF4GI; PARK18; EIF-4G1
SwissProt:	G5E9S1
IHC positive control:	Human lung carcinoma
IHC Recommend dilution:	50-100
WB Predicted band size:	220 kDa
WB Positive control:	HT29 cells treated with Anisomycin
WB Recommended dilution:	500-1000

