

## BCAT1 抗原(重组蛋白)

中文名称: BCAT1 抗原(重组蛋白)

英文名称: BCAT1 Antigen (Recombinant Protein)

别 名: BCT1; PP18; BCATC; ECA39; MECA39; PNAS121

储 存: 冷冻(-20℃)

相关类别: 抗原

概 述:

Fusion protein corresponding to a region derived from 3-301 amino acids of human BCAT1

## 技术规格:

Full name:	branched chain amino acid transaminase 1
Synonyms:	BCT1; PP18; BCATC; ECA39; MECA39; PNAS121
Swissprot:	P54687
Gene Accession:	BC033864
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	This gene encodes the cytosolic form of the enzyme branched-chain amino acid transaminase. This enzyme catalyzes the reversible transamination of branched-chain alpha-keto acids to branched-chain L-amino acids essential for cell growth. Two different clinical disorders have been attributed to a defect of branched-chain amino acid transamination: hypervalinemia and hyperleucine-isoleucinemia. As there is also a gene encoding a mitochondrial form of this enzyme, mutat



ions in either gene may contribute to these disorders. Alternatively spliced transcript variants have been described.