

兔抗 MALT1 多克隆抗体

中文名称：兔抗 MALT1 多克隆抗体

英文名称：Anti-MALT1 rabbit polyclonal antibody

别名：DKFZp434L132; MLT; MLT1

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：MALT1

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

| | |
|--------------------|--|
| Background: | Mucosa-associated lymphoid tissue translocation gene 1 (MALT1) is a paracaspase that is a critical mediator of T-cell receptor activation of NF-κB and may contribute to the progression of MALT lymphomas. It contains two immunoglobulin-like domains, an amino-terminal death domain and a carboxy-terminal caspase-like domain. Association of MALT1 with Bcl-10 and CARD11/Carma1 leads to activation of IKK and subsequent stimulation of NF-κB, resulting in increased proliferation and inhibition of apoptosis. A common translocation in MALT B-cell non-Hodgkin lymphomas t(11;18)(q21;q21) results in the fusion of the amino termin |
|--------------------|--|

| | |
|---------------------------------|--|
| | us of API2 (c-IAP2), a member of the inhibitor of apoptosis protein family, to the carboxy terminus of MALT1. The API2-MALT1 fusion protein likely leads to deregulation of NF- κ B, contributing to increased oncogenic potential. |
| Applications: | WB |
| Name of antibody: | MALT1 |
| Immunogen: | Fusion protein of human MALT1 |
| Full name: | mucosa associated lymphoid tissue lymphoma translocation gene 1 |
| Synonyms : | DKFZp434L132; MLT; MLT1 |
| SwissProt: | Q9UDY8 |
| WB Predicted band size: | 92 kDa |
| WB Positive control: | Jurkat and K562 cells, bone marrow tissue |
| WB Recommended dilution: | 500-2000 |

