

兔抗 MAPK14 (Phospho-Tyr182) 多克隆抗 体

中文名称: 兔抗 MAPK14 (Phospho-Tyr182) 多克隆抗体

英文名称: Anti-MAPK14 (Phospho-Tyr182) rabbit polyclonal antibody

别名: RP1-179N16.5; CSBP; CSBP1; CSBP2; CSPB1; EXIP; Mxi2; PRKM14; PRKM15; RK;

SAPK2A; p3 8; P38alpha

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

抗 原: MAPK14 (Phospho-Tyr182)

反应种属: Human, Mouse, Rat

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:

Responds to activation by environmental stress, pro-inflam matory cytokines and lipopolysaccharide (LPS) by phosphor ylating a number of transcription factors, such as ELK1 and ATF2 and several downstream kinases, such as MAPKAPK2 and MAPKAPK5. Plays a critical role in the production of so me cytokines, for example IL-6. May play a role in stabilization of EPO mRNA during hypoxic stress. Isoform Mxi2 activation is stimulated by mitogens and oxidative stress and o



	nly poorly phosphorylates ELK1 and ATF2. Isoform Exip may play a role in the early onset of apoptosis.
Applications:	WB, IHC
Name of antibody:	MAPK14 (Phospho-Tyr182)
Immunogen:	Synthetic peptide of human MAPK14 (Phospho-Tyr182)
Full name:	mitogen-activated protein kinase 14 (Phospho-Tyr182)
Synonyms :	RP1-179N16.5; CSBP; CSBP1; CSBP2; CSPB1; EXIP; Mxi2; PRK M14; PRKM15; RK; SAPK2A; p38; P38alpha
SwissProt:	Q16539
IHC positive control:	Human breast carcinoma
IHC Recommend dilution:	50-100
WB Predicted band size:	43 kDa
WB Positive control:	Hela cells treated with UV
WB Recommended dilution:	500-1000





