

兔抗 ME3 多克隆抗体

中文名称: 兔抗 ME3 多克隆抗体

英文名称: Anti-ME3 rabbit polyclonal antibody

别 名: NADP-ME

储 存: 冷冻(-20℃) 避光

抗 原: ME3

宿 主: Rabbit

相关类别: 一抗

反应种属: Human Mouse

标 记 物: Unconjugate

克隆类型: Unconjugate

技术规格

| Background: | Malic enzyme catalyzes the oxidative decarboxylation of |
|---------------|--|
| | malate to pyruvate using either NAD+ or NADP+ as a co |
| | factor. Mammalian tissues contain 3 distinct isoforms of |
| | malic enzyme: a cytosolic NADP(+)-dependent isoform, a |
| | mitochondrial NADP(+)-dependent isoform, and a mitocho |
| | ndrial NAD(+)-dependent isoform. This gene encodes a m |
| | itochondrial NADP(+)-dependent isoform. Multiple alternat |
| | ively spliced transcript variants have been found for this g |
| | ene, but the biological validity of some variants has not b |
| | een determined. |
| Applications: | WB |



| Name of antibody: | ME3 |
|--------------------------|---|
| Immunogen: | Synthesized peptide derived from internal of human ME3. |
| Full name: | malic enzyme 3, NADP(+)-dependent, mitochondrial |
| Synonyms: | NADP-ME |
| SwissProt: | Q16798 |
| WB Predicted band size: | 67 kDa |
| WB Positive control: | NIH/3T3 cells and RAW264.7 cells lysates |
| WB Recommended dilution: | 500-3000 |

