

兔抗 H2AJ 多克隆抗体

中文名称：兔抗 H2AJ 多克隆抗体

英文名称： Anti-H2AJ rabbit polyclonal antibody

别名： H2A.J histone; H2AFJ

相关类别： 一抗

储存： 冷冻（-20℃）

宿主： Rabbit

抗原： H2AJ

反应种属： Human, Mouse, Rat

标记物： Unconjugate


克隆类型： rabbit polyclonal



技术规格

Background:

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is located on chromosome 12 and encodes a replication-independent histone that is a variant H

	2A histone. The protein is divergent at the C-terminus compared to the consensus H2A histone family member. This gene also encodes an antimicrobial peptide with antibacterial and antifungal activity.[provided by RefSeq, Oct 2015]
Applications:	ELISA, WB, IHC
Name of antibody:	H2AJ
Immunogen:	Fusion protein of human H2AJ
Full name:	H2A.J histone
Synonyms:	H2AFJ
SwissProt:	Q9BTM1
IHC positive control:	Human esophagus cancer and Human breast cancer
IHC Recommend dilution:	200-300
WB Predicted band size:	14 kDa
WB Positive control:	Jurkat cell lysate
WB Recommended dilution:	500-2000





mlbio 酶联生物
Good elisakit producers