

兔抗 HSD17B4 多克隆抗体

中文名称：兔抗 HSD17B4 多克隆抗体

英文名称：Anti-HSD17B4 rabbit polyclonal antibody

别名：DBP; MFE-2; MPF-2; PRLTS1; SDR8C1

相关类别：一抗

储存：冷冻（-20℃）

抗原：HSD17B4

宿主：Rabbit

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	The protein encoded by this gene is a bifunctional enzyme that is involved in the peroxisomal beta-oxidation pathway for fatty acids. It also acts as a catalyst for the formation of 3-ketoacyl-CoA intermediates from both straight-chain and 2-methyl-branched-chain fatty acids. Defects in this gene that affect the peroxisomal fatty acid beta-oxidation activity are a cause of D-bifunctional protein deficiency (DBPD). An apparent pseudogene of this gene is present on chromosome 8.
Applications:	ELISA, WB, IHC
Name of antibody:	HSD17B4
Immunogen:	Fusion protein of human HSD17B4

Full name:	hydroxysteroid (17-beta) dehydrogenase 4
Synonyms :	DBP; MFE-2; MPF-2; PRLTS1; SDR8C1
SwissProt:	P51659
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human colon cancer and human cervical cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	80 kDa
WB Positive control:	Huvec and K562 cells
WB Recommended dilution:	500-2000



