

## 兔抗 NFATC3(Phospho-Ser165) 多克隆抗体

中文名称：兔抗 NFATC3(Phospho-Ser165) 多克隆抗体

英文名称：Anti-NFATC3(Phospho-Ser165) rabbit polyclonal antibody

别名：NFAT4; NFATX

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：NFATC3(Phospho-Ser165)

反应种属：Human Mouse

标记物：Unconjugate

克隆类型：Unconjugate

### 技术规格

|                          |   |
|--------------------------|---|
| <b>Background:</b>       | The product of this gene is a member of the nuclear factors of activated T cells DNA-binding transcription complex. This complex consists of at least two components: a preexisting cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation and an inducible nuclear component. Other members of this family participate to form this complex also. |
| <b>Applications:</b>     | WB  |
| <b>Name of antibody:</b> | NFATC3(Phospho-Ser165)  |
| <b>Immunogen:</b>        | Peptide sequence around phosphorylation site of Serine 165 (S-P-S(p)-P-A) derived from Human NFAT4.   |

|                                 |   |
|---------------------------------|---|
| <b>Full name:</b>               | nuclear factor of activated T-cells, cytoplasmic, calcineurin-d<br>ependent 3 |
| <b>Synonyms :</b>               | NFAT4; NFATX  |
| <b>SwissProt:</b>               | Q12968  |
| <b>WB Predicted band size:</b>  | 116 kDa   |
| <b>WB Positive control:</b>     | Hela cells lysates  |
| <b>WB Recommended dilution:</b> | 500-1000  |

