

兔抗 NLRP3 多克隆抗体

中文名称: 兔抗 NLRP3 多克隆抗体

英文名称: Anti-NLRP3 rabbit polyclonal antibody

别名: AII; AVP; FCU; MWS; FCAS; CIAS1; NALP3; C1orf7; CLR1.1; PYPAF1; AGTAVPRL

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗原: NLRP3

相关类别: 一抗

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:

This gene encodes a pyrin-like protein containing a pyrin domain, a nucleotide-binding site (NBS) domain, and a leu cine-rich repeat (LRR) motif. This protein interacts with the apoptosis-associated speck-like protein PYCARD/ASC, which contains a caspase recruitment domain, and is a member of the NALP3 inflammasome complex. This complex functions as an upstream activator of NF-kappaB signaling, and it plays a role in the regulation of inflammation, the immune response, and apoptosis. Mutations in this gene are associated with familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome, and



| | neonatal-onset multisystem inflammatory disease (NOMID). Multiple alternatively spliced transcript variants encoding d istinct isoforms have been identified for this gene. Alterna tive 5' UTR structures are suggested by available data; ho wever, insufficient evidence is available to determine if all of the represented 5' UTR splice patterns are biologically valid. |
|-----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Applications: | ELISA, IHC |
| Name of antibody: | NLRP3 |
| Immunogen: | Fusion protein of human NLRP3 |
| Full name: | NLR family, pyrin domain containing 3 |
| Synonyms : | AII; AVP; FCU; MWS; FCAS; CIAS1; NALP3; C1orf7; CLR1.1; PYPAF1; AGTAVPRL |
| SwissProt: | Q96P20 |
| ELISA Recommended dilution: | 1000-2000 |
| IHC positive control: | Human esophagus cancer and human thyroid cancer |
| IHC Recommend dilution: | 15-50 |



