

兔抗 RPL17 多克隆抗体

中文名称：兔抗 RPL17 多克隆抗体

英文名称：Anti-RPL17 rabbit polyclonal antibody

别名：ribosomal protein L17; L17; PD-1; RPL23

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：RPL17

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L22P family of ribosomal proteins. It is located in the cytoplasm. This gene has been referred to as rpl23 because the encoded protein shares amino acid identity with ribosomal protein L23 from <i>Halobacterium marismortui</i> ; however, its official symbol is RPL17. As is typical for genes encodi
--------------------	---

	ng ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. Alternative splicing results in multiple transcript variants. Read-through transcription also exists between this gene and the neighboring downstream C18orf32 (chromosome 18 open reading frame 32) gene.
Applications:	ELISA, WB, IHC
Name of antibody:	RPL17
Immunogen:	Fusion protein of human RPL17
Full name:	ribosomal protein L17
Synonyms:	L17; PD-1; RPL23
SwissProt:	P18621
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human tonsil
IHC Recommend dilution:	50-200
WB Predicted band size:	21 kDa
WB Positive control:	293T cell, Mouse brain tissue, Mouse testis tissue, HepG2 cell, Mouse liver tissue, A375 cell lysates
WB Recommended dilution:	1000-5000



