

兔抗 PLD2(Phospho-Tyr169) 多克隆抗体

中文名称：兔抗 PLD2(Phospho-Tyr169) 多克隆抗体

英文名称：Anti-PLD2(Phospho-Tyr169) rabbit polyclonal antibody

别名：PLD2

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：PLD2(Phospho-Tyr169)

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	The protein encoded by this gene catalyzes the hydrolysis of phosphatidylcholine to phosphatidic acid and choline. The activity of the encoded enzyme is enhanced by phosphatidylinositol 4,5-bisphosphate and ADP-ribosylation factor-1. This protein localizes to the peripheral membrane and may be involved in cytoskeletal organization, cell cycle control, transcriptional regulation, and/or regulated secretion. Two transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jul 2011]
Applications:	WB, IHC
Name of antibody:	PLD2(Phospho-Tyr169)

Immunogen:	Peptide sequence around phosphorylation site of tyrosine 169(E-N-Y(p)-L-N) derived from Human PLD2.
Full name:	phospholipase D2
Synonyms :	PLD2
SwissProt:	O14939
IHC positive control:	Human brain tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	106 kDa
WB Positive control:	Jurkat cells lysates treated with TNF
WB Recommended dilution:	500-1000



