

兔抗 SLC18A2 多克隆抗体

中文名称：兔抗 SLC18A2 多克隆抗体

英文名称：Anti-SLC18A2 rabbit polyclonal antibody

别名：SVAT, SVMT, VAT2, VMAT2

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：SLC18A2

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	The vesicular monoamine transporter acts to accumulate cytosolic monoamines into synaptic vesicles, using the proton gradient maintained across the synaptic vesicular membrane. Its proper function is essential to the correct activity of the monoaminergic systems that have been implicated in several human neuropsychiatric disorders. The transporter is a site of action of important drugs, including reserpine and tetrabenazine .
Applications:	ELISA, WB, IHC
Name of antibody:	SLC18A2

Immunogen:	Synthetic peptide of human SLC18A2
Full name:	solute carrier family 18 (vesicular monoamine), member 2
Synonyms :	SVAT, SVMT, VAT2, VMAT2
SwissProt:	Q05940
ELISA Recommended dilution:	2000-60000
IHC positive control:	Human brain
IHC Recommend dilution:	25-100
WB Predicted band size:	55 kDa
WB Positive control:	Human cervical cancer tissue
WB Recommended dilution:	500-2000



