

兔抗 PPIA 多克隆抗体

中文名称: 兔抗 PPIA 多克隆抗体

英文名称: Anti-PPIA rabbit polyclonal antibody

别 名: CYPA; CYPH; HEL-S-69p

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗原: PPIA

反应种属: Human, Mouse, Rat

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:	This gene encodes a member of the peptidyl-prolyl cistrans isomerase (PPIase) family. PPIases catalyze the cistrans isomerization of proline imidic peptide bonds in o ligopeptides and accelerate the folding of proteins. The encoded protein is a cyclosporin binding-protein and m ay play a role in cyclosporin A-mediated immunosuppre ssion. The protein can also interact with several HIV proteins, including p55 gag, Vpr, and capsid protein, and h as been shown to be necessary for the formation of inf
	ectious HIV virions.
Applications:	ELISA, WB, IHC



Name of antibody:	PPIA
Immunogen:	Fusion protein of human PPIA
Full name:	peptidylprolyl isomerase A (cyclophilin A)
Synonyms:	CYPA; CYPH; HEL-S-69p
SwissProt:	P62937
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human cervical cancer
IHC Recommend dilution:	40-150
WB Predicted band size:	18 kDa
WB Positive control:	PC3, Raji, NIH/3T3, K562, hela, hepg2 and Jurkat cells
WB Recommended dilution:	500-2000





