

## 兔抗 PRKD1/2/3(Phospho-Ser738/742) 多克 隆抗体

中文名称: 兔抗 PRKD1/2/3(Phospho-Ser738/742) 多克隆抗体

英文名称: Anti-PRKD1/2/3(Phospho-Ser738/742) rabbit polyclonal antibody

别名: PKD; PKCM; PRKCM; PKC-MU

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

抗 原: PRKD1/2/3(Phospho-Ser738/742)

反应种属: Human Mouse Rat

标记物: Unconjugate

克隆类型: Unconjugate

## 技术规格

**Background:** 

Serine/threonine-protein kinase that converts transient diac ylglycerol (DAG) signals into prolonged physiological effects downstream of PKC, and is involved in the regulation of M APK8/JNK1 and Ras signaling, Golgi membrane integrity an d trafficking, cell survival through NF-kappa-B activation, cell migration, cell differentiation by mediating HDAC7 nuclea r export, cell proliferation via MAPK1/3 (ERK1/2) signaling, and plays a role in cardiac hypertrophy, VEGFA-induced an



	giogenesis, genotoxic-induced apoptosis and flagellin-stimul ated inflammatory response. Phosphorylates the epidermal growth factor receptor (EGFR) on dual threonine residues, which leads to the suppression of epidermal growth factor (EGF)-induced MAPK8/JNK1 activation and subsequent JUN phosphorylation. Phosphorylates RIN1, inducing RIN1 bindin g to 14-3-3 proteins YWHAB, YWHAE and YWHAZ and increased competition with RAF1 for binding to GTP-bound for m of Ras proteins (NRAS, HRAS and KRAS). Acts downstrea m of the heterotrimeric G-protein beta/gamma-subunit complex to maintain the structural integrity of the Golgi me mbranes, and is required for protein transport along the se cretory pathway. In the trans-Golgi network (TGN), regulate s the fission of transport vesicles that are on their way to the plasma membrane.
Applications:	WB
Name of antibody:	PRKD1/2/3(Phospho-Ser738/742)
Immunogen:	Peptide sequence around phosphorylation site of Serine738 /Serine 742 (E-K-S(p)-F-R-R-S(p)-V-V) derived from Human PKD1/2/3/PKC $\mu$ .
Full name:	protein kinase D1/2/3
Synonyms :	PKD; PKCM; PRKCM; PKC-MU
SwissProt:	Q15139
WB Predicted band size:	102 kDa
WB Positive control:	A549 cells lysates with PMA
WB Recommended dilution:	500-1000



