

## 小鼠抗 HADH 单克隆抗体

中文名称: 小鼠抗 HADH 单克隆抗体

英文名称: Anti-HADH mouse monoclonal antibody

别 名: HAD; HADH1; HADHSC; HCDH; HHF4; MSCHAD; SCHAD

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

宿 主: Mouse

抗 原: HADH

反应种属: Human

标 记 物: Unconjugate

克隆类型: mouse monoclonal

## 技术规格

Background:	This gene is a member of the 3-hydroxyacyl-CoA dehyd
	rogenase gene family. The encoded protein functions in
	the mitochondrial matrix to catalyze the oxidation of st
	raight-chain 3-hydroxyacyl-CoAs as part of the beta-oxi
	dation pathway. Its enzymatic activity is highest with m
	edium-chain-length fatty acids. Mutations in this gene c
	ause one form of familial hyperinsulinemic hypoglycemi
	a. The human genome contains a related pseudogene
	of this gene on chromosome 15. [provided by RefSeq,
	May 2010].
Applications:	WB, IHC



Name of antibody:	HADH
Immunogen:	Fusion protein of human HADH
Full name:	hydroxyacyl-CoA dehydrogenase (HADH), transcript varia nt 2
Synonyms:	HAD; HADH1; HADHSC; HCDH; HHF4; MSCHAD; SCHAD
SwissProt:	Q16836
IHC positive control:	human liver tissue
IHC Recommend dilution:	25-100
WB Predicted band size:	34 kDa
WB Positive control:	HT29 cell lysate
WB Recommended dilution:	500-2000