

## 小鼠抗 HDAC8 单克隆抗体

中文名称: 小鼠抗 HDAC8 单克隆抗体

英文名称: Anti-HDAC8 mouse monoclonal antibody

别 名: histone deacetylase 8; HD8; WTS; RPD3; CDA07; CDLS5; MRXS6; HDACL1

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Mouse

抗 原: HDAC8

反应种属: Human

标 记 物: Unconjugate

克隆类型: mouse monoclonal

## 技术规格

Background:	Histones play a critical role in transcriptional regulation, c ell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein en coded by this gene belongs to class I of the histone dea cetylase family. It catalyzes the deacetylation of lysine res idues in the histone N-terminal tails and represses transcription in large multiprotein complexes with transcriptiona I co-repressors. Multiple transcript variants encoding different isoforms have been found for this gene.
Applications:	WB



Name of antibody:	HDAC8
Immunogen:	Fusion protein of human HDAC8
Full name:	histone deacetylase 8
Synonyms:	HD8; WTS; RPD3; CDA07; CDLS5; MRXS6; HDACL1
SwissProt:	Q9BY41
WB Predicted band size:	42 KD
WB Positive control:	Hela, Molt4, K562, NIH/3T3, 293T and Jurkat cell lysates
WB Recommended dilution:	500-2000