

小鼠抗 STC1 单克隆抗体

中文名称: 小鼠抗 STC1 单克隆抗体

英文名称: Anti-STC1 mouse monoclonal antibody

别 名: stanniocalcin 1; STC

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Mouse

抗 原: STC1

反应种属: Human, Dog

标 记 物: Unconjugate

克隆类型: mouse monoclonal

技术规格

Background:

This gene encodes a secreted, homodimeric glycoprotein th at is expressed in a wide variety of tissues and may have a utocrine or paracrine functions. The gene contains a 5' UTR rich in CAG trinucleotide repeats. The encoded protein cont ains 11 conserved cysteine residues and is phosphorylated by protein kinase C exclusively on its serine residues. The protein may play a role in the regulation of renal and inte stinal calcium and phosphate transport, cell metabolism, or cellular calcium/phosphate homeostasis. Overexpression of human stanniocalcin 1 in mice produces high serum phosp hate levels, dwarfism, and increased metabolic rate. This ge



	ne has altered expression in hepatocellular, ovarian, and br east cancers.
Applications:	WB
Name of antibody:	STC1
Immunogen:	Fusion protein of human STC1
Full name:	stanniocalcin 1
Synonyms:	STC
SwissProt:	P52823
WB Predicted band size:	28 KD
WB Positive control:	MDCK cell lysate
WB Recommended dilution:	500-2000