



小鼠抗 TAPBPL 单克隆抗体

中文名称: 小鼠抗 TAPBPL 单克隆抗体

英文名称: Anti-TAPBPL mouse monoclonal antibody

别 名: TAPBP-R; TAPBPR

储 存: 冷冻 (-20°C) 避光

抗 原: TAPBPL

宿 主: Mouse

反应种属: Human , Dog , Rat , Monkey , Mouse

相关类别: 一抗

标 记 物: Unconjugate

克隆类型: mouse monoclonal

技术规格

Background:	Tapasin, or TAPBP (MIM 601962), is a member of the variable-constant Ig superfamily that links major histocompatibility complex (MHC) class I molecules to the transporter associated with antigen processing (TAP; see MIM 170260) in the endoplasmic reticulum (ER). The TAPBP gene is located near the MHC complex on chromosome 6p21.3. TAPBPL is a member of the Ig superfamily that is localized on chromosome 12p13.3, a region somewhat paralogous to the MHC.[supplied by OMIM, Mar 2008].
Applications:	WB, IHC, IF
Name of antibody:	TAPBPL
Immunogen:	Fusion protein of human TAPBPL



Full name:	TAP binding protein-like (TAPBPL)
Synonyms:	TAPBP-R; TAPBPR
SwissProt:	Q9BX59
IHC positive control:	human pancreas tissue
IHC Recommend dilution:	30-150
WB Predicted band size:	50 kDa
WB Positive control:	HepG2, Hela, SVT2, A549, COS7, Jurkat, MDCK, PC12, MCF-7 cell lysates
WB Recommended dilution:	200-1000