

## 小鼠抗 TXNDC5 单克隆抗体

中文名称: 小鼠抗 TXNDC5 单克隆抗体

英文名称: Anti-TXNDC5 mouse monoclonal antibody

别 名: HCC2; ERP46; HCC-2; STRF8; PDIA15; UNQ364; ENDOPDI

抗 原: TXNDC5

储 存: 冷冻(-20℃) 避光

宿 主: Mouse

反应种属: Human, Monkey

相关类别: 一抗

标 记 物: Unconjugate

克隆类型: mouse monoclonal

## 技术规格

Background:	This gene encodes a protein-disulfide isomerase. Its express ion is induced by hypoxia and its role may be to protect h ypoxic cells from apoptosis. Alternative splicing results in m ultiple transcript variants. Read-through transcription also ex ists between this gene and the neighboring upstream MUT ED (muted homolog) gene. [provided by RefSeq, Dec 2010].
Applications:	WB, IHC
Name of antibody:	TXNDC5
Immunogen:	Fusion protein of human TXNDC5
Full name:	thioredoxin domain containing 5 (TXNDC5), transcript varian t 2
Synonyms:	HCC2; ERP46; HCC-2; STRF8; PDIA15; UNQ364; ENDOPDI
SwissProt:	Q8NBS9



IHC positive control:	carcinoma of human pancreas tissue and human liver tissue ; human pancreas tissue
IHC Recommend dilution:	30-150
WB Predicted band size:	48 kDa
WB Positive control:	HepG2, Hela, A549, COS7, Jurkat, MCF-7 cell lysates
WB Recommended dilution:	500-2000