

## 兔抗 SLC22A18 多克隆抗体

中文名称：兔抗 SLC22A18 多克隆抗体

英文名称：Anti-SLC22A18 rabbit polyclonal antibody

别名：solute carrier family 22 member 18 HET; ITM; BWR1A; IMPT1; TSSC5; ORCTL2; BWSCR1A; SLC22A1L; p45-BWR1A

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：SLC22A18

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

**Background:**

This gene is one of several tumor-suppressing subtransferable fragments located in the imprinted gene domain of 11p15.5, an important tumor-suppressor gene region. Alterations in this region have been associated with the Beckw

	<p>ith-Wiedemann syndrome, Wilms tumor, rhabdomyosarcoma, adrenocortical carcinoma, and lung, ovarian, and breast cancer. This gene is imprinted, with preferential expression from the maternal allele. Mutations in this gene have been found in Wilms' tumor and lung cancer. This protein may act as a transporter of organic cations, and have a role in the transport of chloroquine and quinidine-related compounds in kidney. Several alternatively spliced transcript variants encoding different isoforms have been described.</p>
<b>Applications:</b>	ELISA, IHC
<b>Name of antibody:</b>	SLC22A18
<b>Immunogen:</b>	Synthetic peptide of human SLC22A18
<b>Full name:</b>	solute carrier family 22 member 18
<b>Synonyms:</b>	HET; ITM; BWR1A; IMPT1; TSSC5; ORCTL2; BWSR1A; SLC22A1L; p45-BWR1A
<b>SwissProt:</b>	Q96BI1

<b>ELISA Recommended dilution:</b>	5000-10000
<b>IHC positive control:</b>	Human colorectal cancer;
<b>IHC Recommend dilution:</b>	20-100

